

Light Quark Masses in QCD

Eduardo de Rafael^a *

^aCentre de Physique Théorique, CNRS-Luminy, Case 907, F-13288 Marseille Cedex 9, France

I report on recent work done in collaboration with Laurent Lellouch and Josep Taron [1] concerning the derivation of rigorous lower bounds for the combination of light quark masses $m_s + m_u$ and $m_d + m_u$.

1. INTRODUCTION

The light u , d and s quarks in the QCD Lagrangian have bilinear couplings

$$\mathcal{L}_{\text{QCD}} = \mathcal{L}_{\text{QCD}}^{(0)} - \sum_{q=u,d,s} m_q \bar{q}(x)q(x), \quad (1)$$

with masses $m_q \neq 0$ which explicitly break the chiral symmetry of the Lagrangian. Twenty five years of efforts of many people (including significant contributions from the chairman of this Conference Stephan Narison,) have resulted in the following values [2]:

$$m_u(4\text{GeV}^2) + m_d(4\text{GeV}^2) = (9.5 \pm 1.9)\text{MeV}, \quad (2)$$

and [3,4]

$$m_u(4\text{GeV}^2) + m_s(4\text{GeV}^2) = (149 \pm 25)\text{MeV}. \quad (3)$$

These are values of the running masses at a scale $\mu^2 = 4\text{GeV}^2$ in the $\overline{\text{MS}}$ renormalization scheme. [The choice of scale is of course arbitrary; 4GeV^2 has become the conventional choice of the lattice community. We shall adopt it here as well so as to make easier the comparison of our bounds with their results.] There are quite a few independent determinations, including some from numerical simulations of lattice QCD [5], which now agree within errors with these values.

The values in (2) and (3) are smaller than those obtained in the very early determinations of the light quark masses using QCD sum rules (see e.g. ref. [6] and references therein.) This is partly due to the fact that the experimental determination of $\Lambda_{\overline{\text{MS}}}$ has resulted in larger values than those

quoted in the early 80's, and partly to the appearance of rather large coefficients in the perturbative expansion of the relevant QCD two-point functions.

Since the summer '96 there have appeared some determinations of the light quark masses from new analyses of numerical simulations of lattice QCD [7,8] which find substantially lower values than the ones quoted above. More recently, an independent lattice QCD determination [9] using dynamical Wilson fermions finds results which, for $m_u + m_d$ are in agreement within errors with those of refs. [7,8]; while for m_s they agree rather well, again within errors, with the value quoted in (3). The present situation concerning the values of the light quark masses has thus become rather confusing again, and this is why we have decided to concentrate on *what can be said on rigorous grounds* about the light quark mass values at present, rather than argue in favour or against the reliability of a particular determination.

We shall show that there exist rigorous lower bounds on how small the light quark masses $m_s + m_u$ and $m_d + m_u$ can be. The derivation of the bounds is based on first principles: the fact that two-point functions

$$\Psi(q^2) \equiv i \int d^4x e^{iq \cdot x} \langle 0 | T (\mathcal{O}(x) \mathcal{O}(0)^\dagger) | 0 \rangle \quad (4)$$

of local operators $\mathcal{O}(x)$ obey dispersion relations, and the positivity of the corresponding hadronic spectral function

$$\begin{aligned} \frac{1}{\pi} \text{Im} \Psi(t) &= (2\pi)^3 \sum_{\Gamma} \delta^{(4)}(q - \sum p_{\Gamma}) \times \\ &\times \langle 0 | \mathcal{O}(0) | \Gamma \rangle \langle \Gamma | \mathcal{O}^\dagger(0) | 0 \rangle, \end{aligned} \quad (5)$$

*Invited talk given at the conference "QCD 97", Montpellier, France, July 1997.

where the sum over Γ is extended to all possible on-shell hadronic states with the quantum numbers of the operator $\mathcal{O}$.

We shall be concerned with two types of two-point functions, one where $\mathcal{O}$ is the divergence of the strangeness changing axial current

$$\mathcal{O}(x) \equiv \partial_\mu A^\mu(x) = (m_s + m_u) : \bar{s}(x) i \gamma_5 u(x) :; \quad (6)$$

and another one where $\mathcal{O}$ is the scalar operator $S(x)$, defined as the isosinglet component of the mass term in the QCD Lagrangian

$$\mathcal{O}(x) \equiv S(x) = \hat{m} [\bar{u}(x) u(x) + \bar{d}(x) d(x)], \quad (7)$$

with

$$\hat{m} \equiv \frac{1}{2}(m_u + m_d). \quad (8)$$

We shall refer to the corresponding two-point functions respectively as the “pseudoscalar channel” and the “scalar channel”. Both two-point functions obey dispersion relations which in QCD require two subtractions, and it is therefore appropriate to consider their second derivatives ($Q^2 \equiv -q^2$):

$$\Psi''(Q^2) = \int_0^\infty dt \frac{2}{(t + Q^2)^3} \frac{1}{\pi} \text{Im} \Psi(t). \quad (9)$$

The bounds follow from the restriction of the sum over all possible hadronic states which can contribute to the spectral function to the state(s) with the lowest invariant mass. It turns out that, for the two operators in (6) and (7), these hadronic contributions are well known phenomenologically; either from experiment or from chiral perturbation theory (χ PT) calculations. On the QCD side of the dispersion relation, the two-point functions in question where the quark masses appear as an overall factor, are known in the deep euclidean region where $Q^2 \gg \Lambda_{\text{QCD}}^2$ from perturbative QCD (pQCD) at the four loop level. The leading non-perturbative $\frac{1}{Q^2}$ -power corrections which appear in the operator product expansion when the time ordered product in (4) is evaluated in the physical vacuum [10] are also known. As we shall see, the bounds we find cast serious doubts on the reliability of the recent lattice determinations reported in refs. [7–9].

2. PSEUDOSCALAR CHANNEL

We shall call $\Psi_5(q^2)$ the two-point function in (4) associated to the divergence of the strangeness changing axial current in (6). The lowest hadronic state which contributes to the spectral function in (5) is the K -pole. From eq. (9) we then have

$$\Psi_5''(Q^2) = \frac{2}{(M_K^2 + Q^2)^3} 2f_K^2 M_K^4 + \int_{t_0}^\infty dt \frac{2}{(t + Q^2)^3} \frac{1}{\pi} \text{Im} \Psi_5(t), \quad (10)$$

where $t_0 = (M_K + 2m_\pi)^2$ is the threshold continuum.

It is convenient to introduce the moments $\Sigma_N(Q^2)$ of the hadronic continuum integral

$$\Sigma_N(Q^2) = \int_{t_0}^\infty dt \frac{2}{(t + Q^2)^3} \times \left(\frac{t_0 + Q^2}{t + Q^2} \right)^N \frac{1}{\pi} \text{Im} \Psi_5(t). \quad (11)$$

One is then confronted with a typical moment problem (see e.g. ref. [19].) The positivity of the continuum spectral function $\frac{1}{\pi} \text{Im} \Psi_5(t)$ constrains the moments $\Sigma_N(Q^2)$ and hence the l.h.s. of (10) where the light quark masses appear. The most general constraints among the first three moments for $N = 0, 1, 2$ are:

$$\Sigma_0(Q^2) \geq 0, \quad \Sigma_1(Q^2) \geq 0, \quad \Sigma_2(Q^2) \geq 0; \quad (12)$$

$$\Sigma_0(Q^2) - \Sigma_1(Q^2) \geq 0, \quad \Sigma_1(Q^2) - \Sigma_2(Q^2) \geq 0; \quad (13)$$

$$\Sigma_0(Q^2) \Sigma_2(Q^2) - (\Sigma_1(Q^2))^2 \geq 0. \quad (14)$$

The inequalities in eq. (13) are in fact trivial unless $2Q^2 < t_0$, which constraints the region in Q^2 to too small values for pQCD to be applicable. The other inequalities lead however to interesting bounds which we next discuss.

2.1. Bounds from $\Sigma_0(Q^2) \geq 0$.

The inequality $\Sigma_0(Q^2) \geq 0$ results in a first bound on the running masses:

$$[m_s(Q^2) + m_u(Q^2)]^2 \geq \frac{16\pi^2}{N_c} \frac{2f_K^2 M_K^4}{Q^4} \times \frac{1}{\left(1 + \frac{M_K^2}{Q^2}\right)^3} \frac{1}{\left[1 + \frac{11}{3} \frac{\alpha_s(Q^2)}{\pi} + \dots\right]}. \quad (15)$$

Notice that this bound is non-trivial in the large- N_c limit ($f_K^2 \sim \mathcal{O}(N_c)$) and in the chiral limit ($m_s \sim M_K^2$). The bound is of course a function of the choice of the euclidean Q -value at which the r.h.s. in eq. (15) is evaluated. For the bound to be meaningful, the choice of $Q[\text{GeV}]$ has to be made sufficiently large so that a pQCD evaluation of the QCD factor

$$\mathcal{F}_0^{\text{QCD}}(Q^2) = 1 + \frac{11}{3} \frac{\alpha_s(Q^2)}{\pi} + \dots, \quad (16)$$

associated to the two-point function Ψ_5'' is applicable. The dots in (16) represent higher order terms which have been calculated up to $\mathcal{O}(\alpha_s^3)$ [14], as well as non-perturbative power corrections of $\mathcal{O}\left(\frac{1}{Q^4}\right)$ [6,15], and strange quark mass corrections of $\mathcal{O}\left(\frac{m_s^2}{Q^2}\right)$ and $\mathcal{O}\left(\frac{m_s^4}{Q^4}\right)$ including $\mathcal{O}(\alpha_s)$ terms [16–18,3]. We consider that, $Q \gtrsim 1.4 \text{ GeV}$ is a safe choice for a numerical estimate of the bounds: the evaluation of $\mathcal{F}_0^{\text{QCD}}$ at $Q = 1.4 \text{ GeV}$ with or without inclusion of the $\mathcal{O}(\alpha_s^3)$ terms differs by less than 10%.

The lower bound which follows from eq. (15) for $m_u + m_s$ at a renormalization scale $\mu^2 = 4 \text{ GeV}^2$ results in the solid curves shown in Fig. 1 below. These are the lower bounds obtained by let-




Figure 1. Lower bound in MeV for $[m_s + m_u](4 \text{ GeV}^2)$ versus $Q(\text{GeV})$ from eq. (15).

ting $\Lambda_{\overline{MS}}^{(3)}$ vary [20] between 290 MeV (the upper curve) and 380 MeV (the lower curve) and using $f_K = 113 \text{ MeV}$, $M_K = 493.67 \text{ MeV}$. We take the number of active flavours $n_f = 3$ in

all the numerical analyses of this work. Notice that the hadronic continuum integral in (11) is always larger than the contribution from three light flavours; and indeed, we have checked that using $n_f = 4$ in the QCD expressions leads to even higher bounds. Values of the quark masses below the solid curves in Fig. 1 are forbidden by the bounds. For comparison, the horizontal lines in the figure correspond to the central values obtained by the authors of ref. [7]: their “quenched” result is the horizontal upper line; their “unquenched” result the horizontal lower line.

We wish to emphasize that in Fig. 1 we are comparing what is meant to be a “calculation” of the quark masses –the horizontal lines which are the lattice results of ref. [7]– with a bound which in fact can only be saturated in the limit where $\Sigma_0(Q^2) = 0$. Physically, this limit corresponds to the extreme case where the hadronic spectral function from the continuum of states is totally neglected! What the plots show is that, even in that extreme limit, and for values of Q in the range $1.4 \text{ GeV} \lesssim Q \lesssim 1.8 \text{ GeV}$ the new lattice results are already in difficulty with the bounds.

2.2. Bounds from the Quadratic Inequality

The quadratic inequality in (14) results in improved lower bounds for the quark masses which we show in Fig. 2 below. Like in Fig. 1, these




Figure 2. Lower bound in MeV for $[m_s + m_u](4 \text{ GeV}^2)$ from the quadratic inequality.

are the lower bounds obtained by letting $\Lambda_{\overline{MS}}^{(3)}$ vary [20] between 290 MeV (the upper curve) and 380 MeV (the lower curve). Values of the quark

masses below the solid curves in Fig. 2 are forbidden by the bounds. The horizontal lines in this figure are also the same lattice results as in Fig. 1.

The quadratic bound is saturated for a δ -like spectral function representation of the hadronic continuum of states at an arbitrary position and with an arbitrary weight. This is certainly less restrictive than the extreme limit with the full hadronic continuum neglected, and it is therefore not surprising that the quadratic bound happens to be better than the ones from $\Sigma_N(Q^2)$ for $N = 0, 1$, and 2. Notice that the quadratic bound in Fig. 2 is plotted at higher Q -values than the bound in Fig. 1. The evaluation of the corresponding QCD factor with or without inclusion of the $\mathcal{O}(\alpha_s^3)$ terms differs by less than 10% for $Q^2 \geq 4 \text{ GeV}^2$ and we consider therefore that for the evaluation of the quadratic bound $Q \gtrsim 2 \text{ GeV}$ is already a safe choice. We find that even the quenched lattice results of refs. [7,8] are in difficulty with these bounds.

Similar bounds can be obtained for $m_u + m_d$ when one considers the two-point function associated to the divergence of the axial current

$$\partial_\mu A^\mu(x) = (m_d + m_u) : \bar{d}(x) i \gamma_5 u(x) : . \quad (17)$$

The method to derive the bounds is exactly the same as the one discussed above and therefore we only show, in Fig. 3 below, the results for the corresponding lower bounds which we obtain from the quadratic inequality. We find again that the




Figure 3. Lower bound in MeV for $[m_d + m_u](4 \text{ GeV}^2)$ from the quadratic inequality.

lattice QCD results of refs. [7–9] for $m_u + m_d$ are in serious difficulties with these bounds. Notice that in this case the mass corrections in the QCD two-point function are negligible at the plotted Q -values.

3. SCALAR-ISOSCALAR CHANNEL

The basic difference between the “scalar” and “pseudoscalar” channels is that the lowest state which contributes to the hadronic spectral function in the “scalar” channel is not a pole but the $\pi - \pi$ continuum with $J = 0$ and $I = 0$. This contribution provides a lower bound to the full spectral function:

$$\frac{1}{\pi} \text{Im} \Psi(t) \geq \frac{1}{16\pi^2} \sqrt{1 - \frac{4m_\pi^2}{t}} \times 3 |F(t)|^2 \theta(t - 4m_\pi^2), \quad (18)$$

where $F(t)$ denotes the scalar-isoscalar pion form factor

$$\langle \pi^a(p) \pi^b(p') | S(0) | 0 \rangle = \delta_{ab} F(t), \quad (19)$$

with $t = (p + p')^2$. In particular, the value of this form factor at the origin is the so called *pion sigma term*. This form factor has been well studied in χ PT at the one and two loop level (see e.g. refs[21,22] and references therein), and the predicted low energy shape agrees well with the available experimental information. With

$$F(t) = F(0) \left[1 + \frac{1}{6} \langle r^2 \rangle_s^\pi t + \mathcal{O}(t^2) \right], \quad (20)$$

it is found [22] that

$$F(0) = m_\pi^2 \times (0.99 \pm 0.02) + \mathcal{O}(m_\pi^6), \quad (21)$$

and

$$\langle r^2 \rangle_s^\pi = (0.59 \pm 0.02) \text{ fm}^2. \quad (22)$$

The pion mass used in eq. (21) in ref. [22] is the π^+ -mass. Since the electromagnetic interactions are neglected, we find it more appropriate to use the π^0 -mass instead, which in any case results in a lower contribution to the average quark mass. We also wish to point out that in the generalized version of χ PT (see e.g. ref. [23] and references therein,) the value of $F(0)$ could be sensibly

larger, which would result in even larger lower bounds for the quark masses. A value for the curvature of the scalar form factor is also quoted in ref. [22]; it involves however some extra phenomenological assumptions which go beyond χ PT and this is why we have not included this extra information here.

Using standard methods, (see ref. [24] and references therein, and also refs.[25–27]) one can construct bounds on $\hat{m}$ when something is known about the scalar–isoscalar pion form factor $F(t)$. For this, it is convenient to map the complex q^2 –plane to the complex unit disc:

$$i \frac{1+z}{1-z} = \sqrt{\frac{t}{4m_\pi^2}} - 1. \quad (23)$$

With $F(0)$ and $F'(0)$ as input, the resulting lower bound for $\hat{m}^2$ is

$$\hat{m}^2(Q^2) \geq \frac{3\pi}{N_c} \frac{4m_\pi^2}{Q^4} \frac{F(0)^2}{\left(\sqrt{1 + \frac{4m_\pi^2}{Q^2}} + \frac{2m_\pi}{Q}\right)^6} \times \frac{1 + \left\{-\frac{1}{2} + 3z[-Q^2] - \frac{8m_\pi^2}{3}\langle r^2 \rangle_s\right\}^2}{\left[1 + \frac{11}{3} \frac{\alpha_s(Q^2)}{\pi} + \dots\right]}. \quad (24)$$

Notice that this bound is somewhat similar to the *first bound* obtained from the divergence of the axial current in eq. (15). There are however some interesting differences which we wish to point out. From the point of view of the large- N_c expansion, the r.h.s. of eq. (24) is $1/N_c$ –suppressed. Also, from the point of view of the chiral expansion the r.h.s. of eq. (24) has an extra factor m_π^2 [see eq. (21)] as compared to the r.h.s. of eq. (15). One might expect therefore that the lower bound in (24) should be much worse than the one in (15). Yet the bound for $\hat{m}$ in eq.(24) is surprisingly competitive. The reason for this is twofold. On the one hand, the absence of a prominent narrow resonance in the $J = 0, I = 0$ channel is compensated by a rather large contribution from the chiral loops, a phenomenological feature which appears to be common to all the explored physical processes where the $J = 0, I = 0$ channel contributes. On the other hand the fact that, numerically, m_π^2 and f_π^2 are not so different. The lower bounds for the sum of running

masses $(m_u + m_d)$ at the renormalization scale $[\mu^2 = 4\text{GeV}^2]$ which follow from this *scalar channel bound* are the dashed curves shown in Fig 4 below. The dashed curves in Fig 4 correspond to




Figure 4. *Lower bounds in MeV from the Scalar Channel for $[m_d + m_u](4\text{GeV}^2)$.*

the choice $\Lambda_{\overline{MS}}^{(3)} = 290\text{ MeV}$ (the upper dashed curve) and $\Lambda_{\overline{MS}}^{(3)} = 380\text{ MeV}$ (the lower dashed curve). The plots show that for values of Q in the range $1.4\text{ GeV} \lesssim Q \lesssim 1.6\text{ GeV}$ the lattice QCD results in [7–9] are dangerously close to lower bounds obtained with very little phenomenological input, (just the values of the scalar form factor at the origin and its slope at the origin as well.) Notice also that the relevant two–point function in this channel is practically the same as in the pseudoscalar channel for which $Q \gtrsim 1.4\text{ GeV}$ is already a safe choice.

It is still possible to improve these bounds by taking into account the extra information provided by the fact that the phase of the scalar form factor $F(t)$ in the elastic region $4m_\pi^2 \leq t \leq 16m_\pi^2$ is precisely the $I = 0, J = 0$ $\pi - \pi$ scattering phase–shift and there is information on this phase–shift both from χ PT and from experiment. The technology to incorporate this information is discussed in refs.[25–27]. We have restricted the phase–shift input to a conservative region: $4m_\pi^2 \leq t \leq (500\text{ MeV})^2$, where a resummation of the chiral logarithms calculated at the two–loop level in χ PT [21,22] is certainly expected to be reliable and is in good agreement with experiment. This results in improved bounds which

correspond to the solid curves shown in Fig. 4. The two curves reflect the variation within the quoted errors of the input parameters. Again, we find that the lattice determinations of refs. [7–9] do not satisfy lower bounds which only incorporate very little hadronic input.

4. CONCLUSIONS

We conclude that the lattice results of refs. [7–9] are in serious difficulties with rigorous lower bounds which can be derived from general properties of QCD and with a minimum of well established phenomenological input; and this in two different hadronic channels. The lower bounds we have derived are perfectly compatible with the sum rules results of refs. [2–4] and earlier references therein. They are also compatible with a recent semiempirical determination of m_s made by the ALEPH collaboration [28] based on the rate of the hadronic τ -decays into strange particles observed at LEP.

Other work with various claims on possible values of the light quark masses has recently appeared in refs. [29–32].

REFERENCES

1. L. Lellouch, E. de Rafael and J. Taron, hep-ph/9707523; to appear in Phys. Letters **B**
2. J. Bijmans, J. Prades, and E. de Rafael, Phys. Lett. **348B** (1995) 226; and references therein.
3. M. Jamin and M. Münz, Z. Phys. **C66** (1995) 633; and references therein.
4. K.G. Chetyrkin *et al*, Phys. Rev. **D51** (1995) 5090; and references therein.
5. C.R. Allton *et al*, Nucl. Phys. **B431** (1994) 667.
6. C. Becchi *et al*, Z. Phys. **C8** (1981) 335.
7. R. Gupta and T. Bhattacharya, Phys. Rev. **D55** (1997) 7203.
8. B.J. Gough *et al*, Phys. Rev. Lett. **79** (1997) 1622.
9. N. Eicker *et al*, SESAM–Collaboration, preprint hep-lat/9704019.
10. M.A. Shifman, A.I. Vainshtein, and V.I. Zakharov, Nucl. Phys. **B147** (1979) 385,448.
11. L.R. Surguladze, Sov. J. Nucl. Phys. **50** (1989) 372.
12. S.G. Gorishny *et al*, Mod. Phys. Lett. **A5** (1990) 2703.
13. S.G. Gorishny *et al*, Phys. Rev. **D43** (1991) 1633.
14. K.G. Chetyrkin, Phys. Lett. **B390** (1997) 309.
15. P. Pascual and E. de Rafael, Z. Phys. **C12** (1982) 127.
16. S.C. Generalis, J. Phys. G **16** (1990) 785.
17. V.P. Spiridonov and K.G. Chetyrkin, Sov. J. Nucl. Phys. **47** (1988) 522.
18. K.G. Chetyrkin, S.G. Gorishny, and V.P. Spiridonov, Phys. Lett. **B160** (1985) 149.
19. N.I. Ahiezer and M. Krein, “Some Questions in the Theory of Moments”, Translations of Math. Monographs, Vol. 2; Amer. Math. Soc., Providence, Rhode Island (1962).
20. *Review of Particle Properties*, Phys. Rev. **D54** (1996) 1.
21. J.F. Donoghue, J. Gasser, and H. Leutwyler, Nucl. Phys. **B343** (1990) 341.
22. J. Gasser and U.G. Meissner, Nucl. Phys. **B357** (1991) 90.
23. M. Knecht and J. Stern, *Generalized Chiral perturbation Theory*, in **The Second Daphne Physics Handbook, Vol I** (1995), 169.
24. E. de Rafael and J. Taron, Phys. Rev. **D50** (1994) 373.
25. G. Auberson, G. Mahoux, and F.R.A. Simão, Nucl. Phys. **B98** (1975) 204.
26. G. Auberson *et al*, Ann. Inst. Henri Poincaré **XXII** (1975), 317.
27. M. Micu, Phys. Rev. **D7** (1973) 2136.
28. M. Davier, Nucl. Phys. **B** (Proc. Suppl.) **55C** (1997), 395.
29. P. Colangelo *et al*, hep-ph/9704249
30. C.A. Dominguez, hep-ph/9708234.
31. F. Ynduráin, hep-ph/9708300.
32. H. Dosch and S. Narison, hep-ph/9709215.

Discussions

C. Allton, UKqcd collaboration

Lattice theorists use a renormalization scale of $\mu = 2\text{GeV}$ since they don't believe perturbation theory down to $\mu = 1\text{GeV}$.

E. de Rafael

I am aware of that. I understand of course that pQCD is only valid at sufficiently large euclidean scales; however, once you have pinned down the quark masses at a given scale, you can always use the renormalization group equation to run them down to whatever conventional scale (larger than $\Lambda_{\overline{MS}}$) you want.

A. Pivovarov, Moscow

How worse (or better?) would be your bounds if you switch off QCD corrections at all?

E. de Rafael

As I said, the pQCD corrections are large and positive; therefore, if ignored, the lower bounds for the light quark masses would be significantly larger.